

Lowest order covariant averaging of a perturbed metric and of the Einstein tensor

Dieter Gromes

Institut für Theoretische Physik der Universität Heidelberg
Philosophenweg 16, D-69120 Heidelberg
E - mail: d.gromes@thphys.uni-heidelberg.de

Abstract: We present an explicit averaging formula in lowest order. Besides an arbitrary smearing function it contains two integrals of this function. This is necessary in order to achieve covariance. There is no need to solve any equations. In three dimensions the same averaging formula yields a covariant averaging of the Einstein tensor and thus of the field equations. We also present a simple extension to static perturbations in four dimensions. Various further extensions of the formalism appear possible.

1 Introduction

The averaging problem in general relativity was first raised by Shirkov and Fisher [1] in 1963. The energy-momentum-tensor used in cosmological models is an average over the non homogeneous tensor present in nature. But, due to the non linear nature of Einstein's equations, the metric belonging to the averaged energy-momentum tensor is not identical to the averaged metric. The averaging prescription by itself provides a fundamental problem because of the freedom of choice of coordinates. The authors of [1] suggested to integrate the metric tensor over a four dimensional volume with the familiar factor $\sqrt{-g}$ in the measure. Such an expression is, however, not covariant due to the freedom of performing local transformations. A covariant averaging prescription can be constructed by introducing a bivector $g_{\alpha}^{\beta}(x, x')$ of geodesic parallel displacement, as discussed in the appendix of [2]. This transforms as a vector with respect to coordinate transformations at either x or x' and maps a vector $A_{\beta}(x')$ to $\bar{A}_{\alpha}(x) = g_{\alpha}^{\beta}(x, x')A_{\beta}(x')$, analogously for higher order tensors. An averaging with the help of bivectors was also used in the work of Zalaletdinov [3] where the emphasis was on the commutativity of averaging and covariant differentiation. As remarked by Stoeger, Helmi, and Torres [4] the method of using a covariantly conserved bivector is not applicable to the metric, because the covariant derivative of the metric vanishes. The metric is therefore invariant under this averaging procedure. In the thesis of Behrend [5] the metric is represented by tetrads and the averaging performed over the latter. The tetrads are chosen according to a covariant minimalization prescription.

This is only an extremely brief survey of the literature. For more references as well as the implications for the fitting problem, back reaction, contributions to dark energy, we refer e.g. to the monograph of Krasinski [6] and to the comprehensive recent review of Buchert [7].

Under a covariant averaging process we understand a prescription which has the following properties. Let two observers describe the same physics in different coordinate systems S and S' , with metric tensors $g_{\mu\nu}$ and $g'_{\mu\nu}$. Both of them apply a definite averaging procedure in their respective systems, resulting in the averaged metrics $\langle g_{\mu\nu} \rangle$ and $\langle g'_{\mu\nu} \rangle$, respectively. Then the results have to be connected by the same transformation as the original metric, i. e.

$$\langle g'_{\mu\nu} \rangle = \langle g_{\mu\nu} \rangle'. \quad (1.1)$$

In other words, the operations of averaging and of coordinate transformations have to commute. Furthermore, averaging over a region which is closely located around some point should, of course, reproduce the metric at this point.

In two respects the approach in the present paper is modest. Firstly we assume that a reasonable foliation into space and time has already been performed, so we will essentially concentrate on spatial averaging of a perturbed flat metric, with time kept fixed. At the end we give a simple special generalization to four dimensions. Secondly we will demonstrate covariance only in first order of the perturbation of the metric.

In other respects our approach is ambitious. We give a closed formula for the averaging prescription, there is no need to solve any differential equations or to resort to a background of dust, perfect fluid, or whatsoever. We not only can average over

a given sphere, but may introduce an arbitrary smearing function $f(r)$. This appears more physical than a sharp cutoff at the boundary of the considered sphere. A very important result, at present only derived in three dimensions, is the fact that we can apply the same formula which works for the metric as well to the Einstein tensor. This implies that the Einstein equations for the averaged metric are identical to the averaged equations.

The mapping of the metric $g_{kl}(x')$ to the averaged metric $\langle g_{mn} \rangle(x)$ is represented by a bitensor $K_{mn}^{kl}(x' - x)$ which we will specify in detail. A product of bivectors, as frequently used in the literature, is not sufficient. A mapping with the help of a bitensor was also formulated by Boersma [8], although without going into details.

The paper is organized as follows.

In sect. 2 we introduce the necessary technicalities for the projective coordinates. Technically these are much more convenient than the familiar polar coordinates, in particular for the many partial integrations which we have to perform. In sect. 3 we present the general form of the averaging formula. A central aspect is that, besides the arbitrary smearing function $f(r)$, the integrals $F(r)$ and $G(r)$ over f and f/r appear. This is essential for the proof of covariance in sect. 4. Fixing the remaining freedom of the parameters appropriately, we show in sect. 5 the surprising and highly welcome result, that our formula also yields a covariant averaging of the Einstein tensor. In sect. 6 we discuss a simple four dimensional covariant generalization for static perturbations. The proof for the covariant averaging of the Einstein tensor “almost” goes through also in this case, but fails at the very end. Principal limitations and possible generalizations are discussed in the conclusions.

2 Projective coordinates

For the moment we choose the point, where the averaging is to be performed, as the origin for simplicity. Instead of using standard polar coordinates $x = r \sin \theta \cos \phi$, $y = r \sin \theta \sin \phi$, $z = r \cos \theta$, it is much more convenient for various reasons to use the coordinates $r, u \equiv u^1, v \equiv u^2$ of stereographic projection. They are connected to the polar coordinates by $u = \tan \frac{\theta}{2} \cos \phi$, $v = \tan \frac{\theta}{2} \sin \phi$, and to the cartesian coordinates by

$$u \equiv u^1 = \frac{x}{r+z}, \quad v \equiv u^2 = \frac{y}{r+z}, \quad r = \sqrt{x^2 + y^2 + z^2}, \quad (2.1)$$

or, vice versa,

$$x = \frac{2u}{1+w^2} r, \quad y = \frac{2v}{1+w^2} r, \quad z = \frac{1-w^2}{1+w^2} r, \quad \text{with } w^2 \equiv u^2 + v^2. \quad (2.2)$$

The geometric meaning is simple. $2\sqrt{w^2}$ is the distance from the north pole $(0, 0, 1)$ to the point where the straight line from the south pole $(0, 0, -1)$ through (x, y, z) on the unit sphere cuts the plane $z = 1$. The north pole corresponds to $u = v = 0$, the south pole to $u^2 + v^2 = \infty$. The integration element for the angular averaging is

$$\frac{d\Omega}{4\pi} = \frac{1}{4\pi} \sin \theta d\theta d\phi = \frac{1}{\pi} \frac{du dv}{[1+w^2]^2}. \quad (2.3)$$

The partial derivatives in projective coordinates become

$$\frac{\partial}{\partial x^l} = \frac{x_l}{r} \frac{\partial}{\partial r} + \frac{\partial u^i}{\partial x^l} \frac{\partial}{\partial u^i}, \quad (2.4)$$

with

$$\frac{\partial u^i}{\partial x^l} = \frac{\epsilon_{3lk}\epsilon_{3ij}x_j - \epsilon_{ij}\epsilon_{jlk}(r+z)}{r(r+z)^2} x^k \text{ (for } l = 1, 2), \frac{\partial u^i}{\partial x^3} = -\frac{u^i}{r}. \quad (2.5)$$

Fortunately (2.5) will not be needed explicitly, we will only need two simple obvious properties of the partial derivatives $\partial u^i / \partial x^l$: They are orthogonal to x^l , i.e. $x^l \partial u^i / \partial x^l = 0$, and proportional to $1/r$ for fixed u, v .

The perturbed three dimensional metric in cartesian coordinates is written as $g_{mn}(x) = \delta_{mn} + h_{mn}(x)$. This decomposition is preserved under translations, rotations, and infinitesimal transformations. In the projective coordinates we also split off the flat part and write (indices i, j run from 1 to 2 in (2.6) and refer to u^1, u^2)

$$g_{rr} = 1 + h_{rr}, \quad g_{ri} = \frac{2r}{1+w^2} h_{ri}, \quad g_{ij} = \frac{4r^2}{[1+w^2]^2} [\delta_{ij} + h_{ij}]. \quad (2.6)$$

A corresponding decomposition can be written down for the polar coordinates.

3 General form of the averaging formula

The formula presented below looks somewhat strange at first sight, therefore it is appropriate to motivate it. Originally we tried to transform to a specific system, perform the average there, and transform back to the original one. In this way covariance is achieved. The conditions which essentially fixed the specific system, denoted by primes, were

$$h'_{rr} - h'_{rr}(0) = h'_{r1} - h'_{r1}(0) = h'_{r2} - h'_{r2}(0) = 0. \quad (3.1)$$

To avoid singularities of the transformation between the systems, it was necessary to split off the values at the origin in these conditions. One could next determine the transformation leading from the original to the primed system, perform the average there, and transform back. The fact that we had to split off the values at the origin had the unwanted consequence that part of the metric remained unaveraged. Instead of pursuing this approach further it is, however, more useful to abstract from this original motivation and to concentrate on the structure which was obtained in this way. The important point is the following. After determining the transformation, choosing a smearing function $f(r)$, and performing the average, not only the original function $f(r)$, but also two integrals $F(r)$ and $G(r)$ appear. This will be the key for achieving covariance. So much for the motivation, all the following is independent of it.

In detail, we choose a smearing function $f(r)$ with

$$\int_0^\infty f(r) dr = 1. \quad (3.2)$$

Because we are in three space dimensions one should have $f(r) \sim r^2$ for $r \rightarrow 0$.

We will further need the function $F(r)$, the integral over $f(r)$, as well as the function $G(r)$, the integral over $f(r)/r$, both normalized such that they vanish at infinity:

$$F(r) = - \int_r^\infty f(r') dr', \quad G(r) = - \int_r^\infty \frac{f(r')}{r'} dr'. \quad (3.3)$$

Obviously one has $F(0) = -1$.

The general form of the three dimensional averaging formula now reads

$$\langle g_{mn} \rangle (0) = \int K_{mn}^{kl} g_{kl}(x, y, z) dr \frac{d\Omega}{4\pi}, \quad (3.4)$$

with

$$K_{mn}^{kl} = K_{[f]mn}^{kl} f(r) + K_{[F/r]mn}^{kl} \frac{F(r)}{r} + K_{[G]mn}^{kl} G(r). \quad (3.5)$$

The structure of the tensors (tensors in the sense of linear algebra) $K_{[a]mn}^{kl}$, ($a = f, F/r, G$) is, with constant coefficients $A_{[a]}, \dots, F_{[a]}$,

$$\begin{aligned} K_{[a]mn}^{kl}(x) = & A_{[a]}(\delta_m^k \delta_n^l + \delta_n^k \delta_m^l) + B_{[a]} \delta_{mn} \delta^{kl} + C_{[a]} \delta_{mn} \frac{x^k x^l}{r^2} + D_{[a]} \delta^{kl} \frac{x_m x_n}{r^2} \\ & + E_{[a]} \left[\delta_m^k \frac{x_n x^l}{r^2} + \delta_m^l \frac{x_n x^k}{r^2} + \delta_n^k \frac{x_m x^l}{r^2} + \delta_n^l \frac{x_m x^k}{r^2} \right] + F_{[a]} \frac{x_m x_n x^k x^l}{r^4}. \end{aligned} \quad (3.6)$$

This is the most general tensor structure which is symmetric under the exchange $m \leftrightarrow n$, and under $k \leftrightarrow l$. Naive averaging would correspond to $A_{[f]} = 1/2$ and all the other 17 coefficients $B_{[f]}, \dots, F_{[G]}$ vanishing.

There are some restrictions for the coefficients from the beginning. The function $F(r)/r$ is singular at $r = 0$. The expansion of g_{kl} around $r = 0$ starts with the constant $g_{kl}(0)$. For this term we can perform the angular averaging using

$$\frac{x_m x_n}{r^2} \rightarrow \frac{1}{3} \delta_{mn}, \quad \frac{x_m x_n x^k x^l}{r^4} \rightarrow \frac{1}{15} (\delta_{mn} \delta^{kl} + \delta_m^k \delta_n^l + \delta_n^k \delta_m^l) \quad \text{etc.} \quad (3.7)$$

The tensor $K_{[F/r]mn}^{kl}$ has to vanish after angular integration in order to avoid a singularity at $r = 0$. Collecting the factors in front of $\delta_m^k \delta_n^l + \delta_n^k \delta_m^l$, and those in front of $\delta_{mn} \delta^{kl}$, this leads to the two conditions

$$A_{[F/r]} + \frac{2}{3} E_{[F/r]} + \frac{1}{15} F_{[F/r]} = 0, \quad (3.8)$$

$$B_{[F/r]} + \frac{1}{3} C_{[F/r]} + \frac{1}{3} D_{[F/r]} + \frac{1}{15} F_{[F/r]} = 0. \quad (3.9)$$

There are also restrictions for the coefficients which multiply $f(r)$ and $G(r)$. These will become relevant for proving covariance.

$K_{[f]mn}^{kl}$ is restricted by the transversality condition $x_l K_{[f]mn}^{kl} = 0$, which implies

$$C_{[f]} = -B_{[f]}, \quad E_{[f]} = -A_{[f]}, \quad F_{[f]} = 2A_{[f]} - D_{[f]}. \quad (3.10)$$

Finally, $K_{[G]mn}^{kl}$ contains only terms $\sim x^k x^l$, which means that

$$A_{[G]} = B_{[G]} = D_{[G]} = E_{[G]} = 0. \quad (3.11)$$

Therefore the expressions for $K_{[f]mn}^{kl}$ and $K_{[G]mn}^{kl}$ simplify to

$$\begin{aligned} K_{[f]mn}^{kl} = & A_{[f]} \left[\left(\delta_m^k - \frac{x_m x^k}{r^2} \right) \left(\delta_n^l - \frac{x_n x^l}{r^2} \right) + (m \leftrightarrow n) \right] \\ & + (B_{[f]} \delta_{mn} + D_{[f]} \frac{x_m x_n}{r^2}) \left(\delta^{kl} - \frac{x^k x^l}{r^2} \right), \end{aligned} \quad (3.12)$$

$$K_{[G]mn}^{kl} = K_{[G]mn} \frac{x^k x^l}{r^2}, \text{ with } K_{[G]mn} = C_{[G]} \delta_{mn} + F_{[G]} \frac{x_m x_n}{r^2}. \quad (3.13)$$

Let us now investigate the limit of a smearing function $f(r)$ which is closely localized around $r = 0$. The same then holds for $F(r)$ and $G(r)$. We may thus put $g_{kl}(x, y, z) = g_{kl}(0)$ in (3.4) and take it out in front of the integral. The angular averages can be performed using again (3.7). One is left with the radial integrals $\int_0^\infty f(r) dr = 1$, and $\int_0^\infty 1 \cdot G(r) dr = -\int_0^\infty r \cdot G'(r) dr = -1$. The term with $F(r)/r$ does not enter, because the angular integration vanishes due to the conditions (3.8), (3.9) derived before. The result has to be identical to $g_{mn}(0)$. This leads to two further conditions, derived from comparing the terms with $g_{mn}(0)$ and those with $\delta_{mn} g_j^j(0)$. Making use of the simplifications which arise from the restrictions for the coefficients referring to f and G one obtains

$$\frac{1}{15} [14A_{[f]} - 2D_{[f]} - 2F_{[G]}] = 1, \quad (3.14)$$

$$\frac{1}{15} [2A_{[f]} + 10B_{[f]} + 4D_{[f]} - 5C_{[G]} - F_{[G]}] = 0. \quad (3.15)$$

The considerations above also show that the averaging of a constant (e.g. of δ_{mn}) gives back this constant. This also implies, that (3.4) can as well be applied to the perturbation, i.e. one can replace $g_{mn} \rightarrow h_{mn}$, $g_{kl} \rightarrow h_{kl}$ there.

4 Conditions for covariance

Let us apply an arbitrary infinitesimal transformation $x^k = x'^k + \xi^k$, which leads to a change $\delta g_{kl} = \xi_{k,l} + \xi_{l,k} \rightarrow 2\xi_{k,l}$, when contracted with the symmetrical tensors in (3.6). Because invariance with respect to translations and to rigid rotations around the point of consideration is manifest, one can restrict to transformations which leave the origin fixed, i.e. $\xi_k(0) = 0$, such that

$$\xi_k(x) = \xi_{k,l}(0) x^l + O(r^2). \quad (4.1)$$

Consider the change of the integrand in (3.4). In a first step we transform the terms $\sim f(r)$ and $\sim G(r)$ by partial integration with respect to r , such that all three contributions become $\sim F(r)/r$. Because all the tensors $K_{[a]mn}^{kl}$ in (3.6) depend on u, v only, but are independent of r , this is rather simple.

$$\int_0^\infty f(r) \xi_{k,l} dr = \xi_{k,l}(0) - \int_0^\infty \frac{F(r)}{r} \left(r \frac{\partial}{\partial r} \xi_{k,l} \right) dr, \quad (4.2)$$

where we used $F(0) = -1$ in the boundary term.

In the term with $G(r)$ we only use the factor x^l/r in (3.13), not the x^k/r . Both of them are well defined and independent of r , they only depend on the angles (or on u, v , respectively) which are still fixed here. One can apply the following chain of partial integrations, where we used (4.1) and $F(0) = -1$ in the boundary terms:

$$\begin{aligned} \int_0^\infty G(r) \frac{x^l}{r} \xi_{k,l} dr &= \int_0^\infty G(r) \left(\frac{\partial}{\partial r} \xi_k \right) dr = - \int_0^\infty \frac{f(r)}{r} \xi_k dr \\ &= - \frac{x^l}{r} \xi_{k,l}(0) + \int_0^\infty \frac{F(r)}{r} \left(\frac{\partial}{\partial r} \xi_k - \frac{\xi_k}{r} \right) dr. \end{aligned} \quad (4.3)$$

All together this leads to the integrand (to be averaged over the angles)

$$\begin{aligned} &2(K_{[f]mn}^{kl} - K_{[G]mn} \frac{x^k x^l}{r^2}) \xi_{k,l}(0) \\ &+ 2 \int_0^\infty \frac{F(r)}{r} \left\{ [-K_{[f]mn}^{kl} r \frac{\partial}{\partial r} + K_{[F/r]mn}^{kl}] \xi_{k,l} + K_{[G]mn} \frac{x^k}{r} \left(\frac{\partial}{\partial r} \xi_k - \frac{\xi_k}{r} \right) \right\} dr. \end{aligned} \quad (4.4)$$

All manipulations which involve partial integrations with respect to r have now been performed. We next consider the curly bracket in (4.4) which has to vanish after averaging over u, v . We introduce the projective coordinates and use

$$\xi_{k,l} = \left(\frac{x_l}{r} \frac{\partial}{\partial r} + \frac{\partial u^i}{\partial x^l} \frac{\partial}{\partial u^i} \right) \xi_k. \quad (4.5)$$

From (2.5) we recall that the partial derivatives $\partial u^i / \partial x^l$ are orthogonal to x^l and proportional to $1/r$. Moving $r \partial / \partial r$ to the right of $\partial u^i / \partial x^l$ in the term $-K_{[f]mn}^{kl} (r \partial / \partial r) \xi_{k,l}$ thus gives an extra contribution. The curly bracket in (4.4) becomes

$$\begin{aligned} &\left\{ [-K_{[f]mn}^{kl} \left(\frac{x_l}{r} \frac{\partial}{\partial r} + \frac{\partial u^i}{\partial x^l} \frac{\partial}{\partial u^i} \right) r \frac{\partial}{\partial r} + K_{[f]mn}^{kl} \frac{\partial u^i}{\partial x^l} \frac{\partial}{\partial u^i} \right. \right. \\ &\quad \left. \left. + K_{[F/r]mn}^{kl} \left(\frac{x_l}{r} \frac{\partial}{\partial r} + \frac{\partial u^i}{\partial x^l} \frac{\partial}{\partial u^i} \right) + K_{[G]mn} \frac{x^k}{r} \left(\frac{\partial}{\partial r} - \frac{1}{r} \right) \right\} \xi_k. \end{aligned} \quad (4.6)$$

The first term contains a second derivative with respect to r . This term has no chance to cancel against anything else, but it vanishes due to the transversality condition $x_l K_{[f]mn}^{kl} = 0$. We are left with terms $\sim \partial / \partial r$ and those $\sim 1/r$ (recall that $\partial u^i / \partial x^l \sim 1/r$). Both of them have to vanish after angular averaging for any ξ_k . Thus we multiply by the integration element (2.3) and remove all partial derivatives $\partial / \partial u^i$ acting on ξ_k by partial integrations. This gives two conditions which arise from collecting terms $\sim \partial / \partial r$ and terms $\sim 1/r$:

$$\frac{\partial}{\partial u^i} \left(\frac{K_{[f]mn}^{kl}}{[1+w^2]^2} r \frac{\partial u^i}{\partial x^l} \right) + \frac{K_{[F/r]mn}^{kl}}{[1+w^2]^2} \frac{x_l}{r} + \frac{K_{[G]mn}}{[1+w^2]^2} \frac{x^k}{r} = 0, \quad (4.7)$$

$$-\frac{\partial}{\partial u^i} \left(\frac{K_{[f]mn}^{kl} + K_{[F/r]mn}^{kl}}{[1+w^2]^2} r \frac{\partial u^i}{\partial x^l} \right) - \frac{K_{[G]mn}}{[1+w^2]^2} \frac{x^k}{r} = 0. \quad (4.8)$$

The sum gives a condition for $K_{[F/r]mn}^{kl}$ alone,

$$-\frac{\partial}{\partial u^i} \left(\frac{K_{[F/r]mn}^{kl}}{[1+w^2]^2} r \frac{\partial u^i}{\partial x^l} \right) + \frac{K_{[F/r]mn}^{kl}}{[1+w^2]^2} \frac{x_l}{r} = 0, \quad (4.9)$$

therefore we will discuss (4.9) and (4.7) in the following.

One now has to use the properties of the partial derivatives $\partial u^i / \partial x^l$ in (2.5), and to introduce the expressions (3.6), (3.12), (3.13) for the $K_{[a]mn}^{kl}$. The further treatment can be greatly simplified by making use of the fact that, from rotation invariance and symmetry, the terms in (4.7) - (4.9) must be a superposition of the form $a(r)\delta_{mn}x^k + b(r)(\delta_m^k x_n + \delta_n^k x_m) + c(r)x_m x_n x^k / r^2$. Therefore there are only three invariants which have to vanish, and we are free to choose simple special cases in order to determine them.

The first choice is to take the trace $m = n$ and put $k = 3$. This gives $[3a(r) + 2b(r) + c(r)]z$. Next one can put $m = n = k = 3$ which gives $[a(r) + 2b(r) + c(r)z^2/r^2]z$. The vanishing of these two expressions implies already the three equations $a(r) = b(r) = c(r) = 0$. While the first choice gives only one relation, the second one gives two relations from collecting the terms $\sim (1 - w^2)/[1 + w^2]^3$ and $\sim (1 - w^2)^3/[1 + w^2]^5$. The three relations such obtained from (4.9) read

$$6A_{[F/r]} + 9B_{[F/r]} + 3C_{[F/r]} + 3D_{[F/r]} + 4E_{[F/r]} + F_{[F/r]} = 0, \quad (4.10)$$

$$6A_{[F/r]} + 3B_{[F/r]} + C_{[F/r]} - 2D_{[F/r]} - 2E_{[F/r]} = 0, \quad (4.11)$$

$$5D_{[F/r]} + 10E_{[F/r]} + F_{[F/r]} = 0. \quad (4.12)$$

The earlier equations (3.8), (3.9) are consequences of (4.10) - (4.12), therefore we can forget them. The same procedure can be applied to (4.7) and leads to three further independent conditions:

$$4A_{[f]} + 6B_{[f]} + 2D_{[f]} - 2A_{[F/r]} - 3B_{[F/r]} - 3C_{[F/r]} - D_{[F/r]} - 4E_{[F/r]} - F_{[F/r]} - 3C_{[G]} - F_{[G]} = 0, \quad (4.13)$$

$$8A_{[f]} + 2B_{[f]} - 2D_{[f]} - 2A_{[F/r]} - B_{[F/r]} - C_{[F/r]} - 2E_{[F/r]} - C_{[G]} = 0, \quad (4.14)$$

$$8A_{[f]} - 4D_{[f]} + D_{[F/r]} + 2E_{[F/r]} + F_{[F/r]} + F_{[G]} = 0. \quad (4.15)$$

If (4.10) - (4.15) are fulfilled, the integral in (4.4), when averaged over the angles, respectively over u, v , vanishes, i.e. the expression is gauge invariant.

Finally we have to consider the boundary term $2(K_{[f]mn}^{kl} - K_{[G]mn}^{kl} x^k x^l / r^2) \xi_{k,l}(0)$ in (4.4). Because $\xi_{k,l}(0)$ is constant, the angular averaging can be performed explicitly. In order to fulfill the covariance condition (1.1), the result must be identical to the change of

$\langle g_{mn} \rangle (0)$ on the lhs, i.e. to $\xi_{m,n}(0) + \xi_{n,m}(0)$. This results in two further conditions which are identical with (3.14), (3.15).

It is worthwhile to mention that the covariance conditions (4.10) - (4.15) were fulfilled in our original approach which was mentioned in the motivation at the beginning of sect. 3. On the other hand, (3.14), (3.15) failed. This failure is due to the fact that there remained contributions which were not averaged.

We have found an averaging formula which is covariant. It contains 11 constant parameters $A_{[f]}, \dots, F_{[G]}$ (3 multiplying $f(r)$, 6 multiplying $F(r)/r$, and 2 multiplying $G(r)$), and has to fulfill 8 independent conditions (3.14), (3.15), (4.10) - (4.15). There is still some freedom which one can use. For reasons to become clear in the next section we impose three further conditions,

$$C_{[a]} = D_{[a]}, \text{ for } a = f, F/r, G. \quad (4.16)$$

This implies a further symmetry of K_{mn}^{kl} :

$$K_{mnkl} = K_{klmn}. \quad (4.17)$$

The terms with A, B, E, F obviously respect this symmetry automatically.

The conditions now fix the parameters uniquely. The result is

	A	B	C	D	E	F
f	3/16	-21/16	21/16	21/16	-3/16	-15/16
F/r	-21/16	27/16	-63/16	-63/16	9/16	225/16
G	0	0	0	0	0	-15/2

(4.18)

5 Covariant averaging of the Einstein tensor

Besides the metric, the Einstein tensor is the most important object in general relativity, because it enters, together with the energy momentum tensor, directly the field equations. It would be highly desirable if one could average the Einstein tensor in exactly the same way as the metric tensor, and if the averaged Einstein tensor would be identical to the Einstein tensor derived from the averaged metric. The old problem that the averaged equations are not identical to the equations with the averaged metric would then disappear.

Let us thus investigate the Einstein tensor. In first order of the perturbation one has

$$2G_{mn} = h_{i,mn}^i + h_{mn,i}^i - h_{m,ni}^i - h_{n,mi}^i - h_{i,j}^i \delta_{mn} + h_{ij}^{,ij} \delta_{mn}. \quad (5.1)$$

Indices are raised and lowered with δ_{ij} here, so their position is in fact irrelevant. The averaging formula (3.4) is now used for an arbitrary point x , the integration variables are denoted by a prime, and $K_{mn}^{kl}(x' - x)$ depends on the difference $x' - x$. The distance r' now means $r' = |x' - x|$. It is convenient to introduce the modified expression

$$\tilde{K}_{mn}^{kl}(x' - x) \equiv \frac{1}{4\pi r'^2} K_{mn}^{kl}(x' - x), \quad (5.2)$$

such that

$$\int K_{mn}^{kl}(x' - x)h_{kl}(x')dr'\frac{d\Omega'}{4\pi} = \int \tilde{K}_{mn}^{kl}(x' - x)h_{kl}(x')d^3x'. \quad (5.3)$$

This makes partial integrations easy in the case under consideration. Differential operators $\partial/\partial x'^n$ acting on $h_{kl}(x')$ can be shifted to $\tilde{K}_{ij}^{kl}(x' - x)$ by partial integration, and finally be replaced by $\partial/\partial x^n$ acting on $\tilde{K}_{ij}^{kl}(x' - x)$.

There are now two possibilities of averaging:

The first possibility is to average the metric in the way described before. Subsequently one calculates the Einstein tensor from (5.1), using the averaged metric on the rhs. The factor of h_{kl} in the integrand then becomes

$$I_{mn}^{kl} = \tilde{K}_i^{ikl},_{mn} + \tilde{K}_{mn,i}^{kl} - \tilde{K}_m^{ikl},_{ni} - \tilde{K}_n^{ikl},_{mi} - \tilde{K}_i^{ikl},_{j}^j \delta_{mn} + \tilde{K}_{ij}^{kl},^{ij} \delta_{mn}. \quad (5.4)$$

The second possibility is to calculate the Einstein tensor G_{kl} within the old metric and then average it with our formula in exactly the same way as we averaged the metric tensor. Shift the partial derivatives from the metric to $\tilde{K}$, and rename dummy indices where necessary such that h_{kl} appears in all six terms. The factor of h_{kl} in the integrand now becomes

$$J_{mn}^{kl} = \tilde{K}_{mn,ij}^{ij} \delta^{kl} + \tilde{K}_{mn,i}^{kl} - \tilde{K}_{mn,i}^{ki} - \tilde{K}_{mn,i}^{li} - \tilde{K}_{mn,i}^{ji} \delta^{kl} + \tilde{K}_{mni}^{i},^{kl}. \quad (5.5)$$

We have to check whether the two expressions are identical. This would be an extremely complicated task if attacked by brute force. Fortunately one can simplify the problem a little bit, although it stays complicated. An inspection of (5.4), (5.5) shows the following property. The terms $\tilde{K}_{mn,i}^{kl}$ are identical. We next use that the tensor $\tilde{K}_{mnkl}$ is invariant under the exchange $(mn) \leftrightarrow (kl)$. The remaining five terms, if arranged properly (e.g. the first in (5.4) and the last in (5.5)), correspond to each other by using this symmetry. This implies $J_{mnkl} = I_{klmn}$. Therefore the difference $I_{mnkl} - J_{mnkl} = I_{mnkl} - I_{klmn}$ is antisymmetric under the exchange $(mn) \leftrightarrow (kl)$. On the other hand, the tensor composition of this expression must have the general form (3.6). The terms $\sim A, B, E, F$ are symmetric under the exchange $(mn) \leftrightarrow (kl)$ and thus cannot appear, therefore the difference has to be of the form

$$I_{mnkl} - J_{mnkl} = c(r')[\delta_{mn} \frac{(x' - x)_k(x' - x)_l}{r'^2} - \delta_{kl} \frac{(x' - x)_m(x' - x)_n}{r'^2}]. \quad (5.6)$$

The knowledge of this structure allows a considerable simplification, because one may now, e.g. contract $k = l$ in order to extract the function $c(r')$. One cannot further contract $m = n$ because then the rhs of (5.6) vanishes identically. Instead of the four indices present originally, one thus has to deal with two indices only and can investigate the expression

$$I_{mnk}^k - J_{mnk}^k = c(r')[\delta_{mn} - 3 \frac{(x' - x)_m(x' - x)_n}{r'^2}]. \quad (5.7)$$

It is now necessary to insert the explicit form of $\tilde{K}_{mn}^{kl}(x' - x)$, to perform the differentiations, and to make use of the relations between the functions $f(r), F(r)/r, G(r)$. The

elementary but tedious calculation gives the structure (5.7) with $c(r)$ a superposition of five terms which are proportional to $f''(r)$, $f'(r)/r$, $f(r)/r^2$, $F(r)/r^3$, $G(r)/r^2$.

Let us, for the moment, keep the parameters free and only make use of the special forms of the tensors $K_{[f]mn}^{kl}$ and $K_{[G]mn}^{kl}$ in (3.12), (3.13), as well, of course, of the conditions $C_{[a]} = D_{[a]}$ in (4.16). The result is striking, therefore we show the explicit result here:

$$\begin{aligned}
c(r) = & 0 \cdot f''(r) \\
& + [2A_{[f]} + 2B_{[f]} - B_{[F/r]} - C_{[F/r]}] f'(r)/r \\
& + [-4B_{[f]} + 7B_{[F/r]} + 6C_{[F/r]} + F_{[F/r]} + F_{[G]}] f(r)/r^2 \\
& + [-15B_{[F/r]} - 10C_{[F/r]} - F_{[F/r]}] F(r)/r^3 \\
& + 0 \cdot G(r)/r^2.
\end{aligned} \tag{5.8}$$

A sort of miracle happens. All five coefficients in front of the functions vanish for the parameters in (4.18)! This implies that the averaged Einstein tensor is identical to the Einstein tensor of the averaged metric, i.e. the averaged field equations are identical to the field equations of the averaged metric.

This property is highly welcome and it is hard to believe that it is accidental. Clearly the symmetry relations shared by our averaging formula and by the Einstein tensor played a central role in the derivation of this result. This becomes evident if one writes

$$2G_{mn} = T_{mn}^{kl} h_{kl}, \tag{5.9}$$

with the operator

$$\begin{aligned}
T_{mn}^{kl} = & \delta^{kl} \partial_m \partial_n + \frac{1}{2} (\delta_m^k \delta_n^l + \delta_n^k \delta_m^l) \partial^i \partial_i - \frac{1}{2} (\delta_m^k \partial_n \partial^l + \delta_m^l \partial_n \partial^k + \delta_n^k \partial_m \partial^l + \delta_n^l \partial_m \partial^k) \\
& - \delta_{mn} \delta^{kl} \partial^j \partial_j + \delta_{mn} \partial^k \partial^l.
\end{aligned} \tag{5.10}$$

Both expressions, $\tilde{K}_{mnkl}$ as well as T_{mnkl} , are symmetric under $m \leftrightarrow n$, under $k \leftrightarrow l$, and under $(m, n) \leftrightarrow (k, l)$. These symmetries implied the vanishing of the five symmetric tensors in the difference of (5.4) and (5.5). The vanishing of the remaining antisymmetric structure in (5.6) could be demonstrated explicitly, but at present we are not aware of some deeper reason behind this.

The result for the covariant averaging of the Einstein tensor is certainly not trivial. For the Ricci tensor, which does not fulfill the above symmetry properties, the relation is not valid.

6 Static perturbations in Minkowski space

Our extension to the four dimensional case is rather modest. We assume that one can find a system in which the perturbation is approximately static. We also neglect the slow time dependence in the Robertson Walker metric, so one may choose coordinates such that the unperturbed metric is the Minkowski one, $(-1, 1, 1, 1)$. To keep this situation,

only rigid translations, rigid spatial rotations, and infinitesimal transformations which keep the time unchanged are allowed. This means that $\xi^0 = 0$, and ξ^m is independent of t . Furthermore we can drop all time derivatives in the metric. Under these restrictions the perturbations h_{00} and h_{m0} become gauge invariant.

We average the perturbation with the following simple ansatz.

$$\langle h_{mn} \rangle (0) = \int \tilde{K}_{mn}^{kl} h_{kl} r^2 dr \frac{d\Omega}{4\pi}, \quad (6.1)$$

$$\langle h_{00} \rangle (0) = \int p(r) h_{00} r^2 dr \frac{d\Omega}{4\pi}, \quad (6.2)$$

$$\langle h_{m0} \rangle (0) = \int q(r) h_{m0} r^2 dr \frac{d\Omega}{4\pi}. \quad (6.3)$$

Of course the functions $p(r)$ and $q(r)$ have to fulfill the normalization conditions

$$\int_0^\infty p(r) r^2 dr = \int_0^\infty q(r) r^2 dr = 1. \quad (6.4)$$

The equations (6.1) - (6.3) are written in such a way that the volume element $d^3x/4\pi$ appears in the integrals. Hopefully, the fact that we sometimes include the factor r^2 of the volume element into the function (as in $f(r)$), and sometimes don't (as in $p(r), q(r)$), does not produce too much confusion. The choice is motivated by the way how we have to perform the various partial integrations.

Equations (6.1) - (6.3) are a simple generalization of our previous formula. For static perturbations they are covariant in the sense of (1.1) with respect to static transformations. One could use a more general ansatz, where a term with h_0^0 is inserted into the rhs of the averaging formula (6.1), and a gauge invariant combination of h_k^k and $(x^k x^l / r^2) h_{kl}$ into (6.2). We found that this does not help to solve the problem which will arise at the end, therefore we keep things simple and work with (6.1) - (6.3).

Let us now consider the (lowest order) Einstein tensor which reads

$$2G_{mn} = 2G_{mn}^{(s)} + h_{0,mn}^0 - h_{0,i}^0 \delta_{mn}, \quad (6.5)$$

$$2G_{00} = h_{i,j}^i - h_{ij}^{,ij}, \quad (6.6)$$

$$2G_{m0} = h_{m0,i}^i - h_{i0,m}^i. \quad (6.7)$$

Here $G_{mn}^{(s)}$ is the spatial part of the Einstein tensor in (5.1). Under the assumptions above, the additional terms in G_{mn} , as well as G_{00} and G_{m0} are invariant under infinitesimal static transformations.

We now investigate whether the Einstein tensor of the averaged metric can be identical to the averaged Einstein tensor. For G_{m0} this is trivial, we start with G_{mn} . For the part $G_{mn}^{(s)}$ we know from the previous section that the result is independent of the order of averaging. We can restrict to the additional contributions in (6.5).

If we average the perturbation h_0^0 according to (6.2) and introduce into (6.5) we obtain the integrand

$$[p_{,mn} - p_{,i}^i \delta_{mn}] h_0^0 = [(p'' - \frac{p'}{r}) \frac{x_m x_n}{r^2} - (p'' + \frac{p'}{r}) \delta_{mn}] h_0^0. \quad (6.8)$$

If, alternatively, we first calculate $2G_{mn}$ in (6.5) with the old metric and then average it in the same way as (6.1), i.e. replace h_{kl} by G_{kl} there on the rhs, and shift the partial derivatives from the perturbation to the multiplying functions, we obtain the integrand

$$[\tilde{K}_{mn,kl}^{kl} - \tilde{K}_{mnk,l}^k] h_0^0. \quad (6.9)$$

This gives the condition

$$\tilde{K}_{mn,kl}^{kl} - \tilde{K}_{mnk,l}^k = (p'' - \frac{p'}{r}) \frac{x_m x_n}{r^2} - (p'' + \frac{p'}{r}) \delta_{mn}. \quad (6.10)$$

We next apply the same procedure to G_{00} . If we average the perturbations h_i^i and h_{ij} according to (6.1) and introduce into (6.6) we obtain the integrand

$$[\tilde{K}_i^{ikl,j} - \tilde{K}_{ij}^{kl,ij}] h_{kl}. \quad (6.11)$$

If, alternatively, we first calculate $2G_{00}$ in (6.6) with the old metric and then average it in the same way as (6.2), i.e. replace h_{00} by G_{00} there, and shift the partial derivatives from the perturbation to the multiplying functions, we obtain the integrand

$$[p_{,j}^j \delta^{kl} - p_{,kl}] h_{kl} = -[(p'' - \frac{p'}{r}) \frac{x^k x^l}{r^2} - (p'' + \frac{p'}{r}) \delta^{kl}] h_{kl}. \quad (6.12)$$

The condition $\tilde{K}_i^{ikl,j} - \tilde{K}_{ij}^{kl,ij} = -[(p'' - \frac{p'}{r}) \frac{x^k x^l}{r^2} - (p'' + \frac{p'}{r}) \delta^{kl}]$ which arises from (6.11), (6.12) is a consequence of (6.10) if one renames the dummy indices k, l there, subsequently replaces m, n by k, l , and uses the symmetry $\tilde{K}_{mnkl} = \tilde{K}_{klmn}$. We thus only need to consider (6.10) in the following.

With (3.5), (3.6) and the parameters in (4.18), a lengthy but elementary calculation gives

$$\tilde{K}_{mn,kl}^{kl} - \tilde{K}_{mnk,l}^k = K_1(r) \frac{x_m x_n}{r^2} + K_2(r) \delta_{mn}, \quad (6.13)$$

with

$$K_1(r) = -\frac{9}{4} \frac{f''(r)}{r^2} + \frac{15}{2} \frac{f'(r)}{r^3} + \frac{9}{2} \frac{f(r)}{r^4} - 30 \frac{G(r)}{r^4}, \quad (6.14)$$

$$K_2(r) = \frac{9}{4} \frac{f''(r)}{r^2} - 3 \frac{f'(r)}{r^3} - 6 \frac{f(r)}{r^4} + 15 \frac{G(r)}{r^4}. \quad (6.15)$$

If one equates the factors of $x_m x_n / r^2$ and δ_{mn} in (6.10) one obtains the two conditions

$$K_1(r) = p''(r) - p'(r)/r, \quad (6.16)$$

$$K_2(r) = -p''(r) - p'(r)/r. \quad (6.17)$$

The sum of these equations gives

$$p'(r) = -\frac{r}{2}(K_1(r) + K_2(r)). \quad (6.18)$$

If one calculates $p''(r)$ from this, and inserts again into (6.16), (6.17) one obtains the integrability condition

$$K_1'(r) + K_2'(r) + \frac{2K_1(r)}{r} = 0. \quad (6.19)$$

This condition consists, in fact, of six conditions, i.e. all the constants in front of $f'''(r)/r^2, f''(r)/r^3, f'(r)/r^4, f(r)/r^5, F(r)/r^6, G(r)/r^5$ have to vanish. Again a sort of miracle occurs. All these constants vanish for the parameters in (4.18).

Up to now everything worked perfectly, but now we run in trouble with the normalization condition (6.4). From (6.18) one gets

$$\begin{aligned} 1 &\stackrel{!}{=} \int_0^\infty r^2 p(r) dr = \left[\frac{r^3}{3} p(r) \right]_0^\infty - \frac{1}{3} \int_0^\infty r^3 p'(r) dr \\ &= \frac{1}{6} \int_0^\infty r^4 (K_1(r) + K_2(r)) dr = \frac{3}{2}. \end{aligned} \quad (6.20)$$

Unfortunately the normalization becomes 3/2 and not 1. The boundary values $[r^3 p(r)]_0^\infty$ cannot help, because these have to vanish in order that $\int_0^\infty r^2 p(r) dr$ converges.

Thus we “almost” succeeded to extend the applicability of the covariant averaging formulae (6.1) - (6.3) to the Einstein tensor, but we failed at the end. There is a clash between the condition from covariance and the normalization. A slightly more general approach can probably solve this problem.

7 Outlook and conclusions

The covariant averaging procedure presented in this paper is complicated. This was to be anticipated, one could not expect to obtain a simple solution to a complicated problem.

Compared to most approaches in the literature our formula shows the following features. First one may be surprised that there is no factor $\sqrt{g}$ in the integrand. We have no comment on this, it simply is not present. An important point is, that it is not sufficient to work with bivectors which mix the indices. It is necessary to have bitensors in (3.6). Our formula is more general than prescriptions which only average over a certain volume because we can use an arbitrary normalized smearing function $f(r)$. But it is essential that, besides the function $f(r)$, also the integrals $F(r)$ and $G(r)$ appear. Two simple examples suggest themselves. We recall that $f(r)$ should behave $\sim r^2$ for small r .

Averaging over a sphere of radius r_0 :

$$f(r) = \frac{3r^2}{r_0^3} \Theta(r_0 - r), \quad F(r) = \left(-1 + \frac{r^3}{r_0^3}\right) \Theta(r_0 - r), \quad G(r) = \left(-\frac{3}{2r_0} + \frac{3r^2}{2r_0^3}\right) \Theta(r_0 - r). \quad (7.1)$$

Averaging with an exponentially decreasing function:

$$f(r) = \frac{r^2}{2r_0^3} e^{-r/r_0}, \quad F(r) = -\left(1 + \frac{r}{r_0} + \frac{r^2}{2r_0^2}\right) e^{-r/r_0}, \quad G(r) = -\left(\frac{1}{2r_0} + \frac{r}{2r_0^2}\right) e^{-r/r_0}. \quad (7.2)$$

One has to face the fact that there is no chance to find a much simpler covariant averaging formula in three dimensions than the one presented here. If one works in first order of the perturbed metric, there must be a linear connection between the original perturbation $h_{kl}(x)$ and the averaged $\langle h_{mn} \rangle(0)$, represented by a tensor (tensor in the sense of linear algebra) K_{mn}^{kl} which is symmetric with respect to $m \leftrightarrow n$ and to $k \leftrightarrow l$. Such a connection has also been discussed by Boersma [8]. The only objects which are available for the construction of K_{mn}^{kl} are the vector x and Kronecker deltas. Therefore one ends up with the six tensors defined in (3.6). The functions in front of the tensors must depend on r only. They are related by covariance. A special solution has been presented in this paper. There are also more general solutions not mentioned here, e.g. averaging formulae which, besides $f(r), F(r)/r, G(r)$, also contain the derivative $rf'(r)$. These may allow to find a prescription which is also suited for a covariant averaging of the Einstein tensor in four dimensions (for static perturbations). The most general form of a covariant averaging formula is under investigation.

A feature of our averaging formula, which might be considered as unpleasant, is the appearance of the integrals $F(r)/r$ and $G(r)$, with $F(0) = -1$ and $G(0)$ finite. Although the term with $F(r)/r$ does not cause a singularity at the origin because the angular integration vanishes there, it somehow hampers the smoothing procedure. Before trying to find a “better” solution one should, however, take notice of the following fact. It is inevitable that functions appear which are not too smooth at the origin. This should be clear from our proof of covariance in sect. 4. An infinitesimal transformation within the averaging formula must result in the corresponding transformation of the averaged metric at the origin. Technically this can only arise through boundary terms at zero which originate from partial integrations. Something substantial must be present near zero in order to produce these boundary terms.

An obvious task to be done is to investigate iterations of our averaging formula and to check whether the iteration procedure converges. But it is clear that one will not always obtain a smooth metric in the limit. The reason is again covariance. We are still free to perform gauge transformations, and by an unfavorable choice of gauge the “smoothed” metric can look wavy and irregular. All one can expect is that the final metric becomes equivalent to a smooth metric.

One could also proceed to extend the approach to second order in the perturbation. We recall that (3.4) also holds if g_{mn} and g_{kl} are replaced by h_{mn} and h_{kl} . In second order an additional contribution which is quadratic in the perturbation will probably be needed in the integral. Though certainly tedious, such an extension appears feasible.

A particularly pleasant property of our formula is the fact that it likewise yields a covariant averaging of the Einstein tensor and thus of the energy momentum tensor in three dimensions, an extra bonus which, quite surprisingly, came out from the suggested averaging formula. We hope to resolve the minor problems found in the four dimensional case by a slightly more general prescription.

Acknowledgement: I thank Juliane Behrend for valuable discussions and for her interest in this work.

References

- [1] M. F. Shirokov, I. Z. Fisher, Sov. Astron. A. J. **6**, 699 (1963), reprinted in Gen. Rel. Grav. **30**, 1411 (1998).
- [2] R. A. Isaacson, Phys. Rev. **166**, 1272 (1968).
- [3] R. M. Zalaletdinov, Gen. Rel. Grav. **24**, 1015 (1992), **25**, 673 (1993). (1964).
- [4] W. R. Stoeger, A. Helmi, D. F. Torres, Int. J. Mod. Phys. **D16**, 1001 (2007), gr-qc/9904020.
- [5] J. Behrend, gr-qc/0812.2859.
- [6] A. Krasinski, Inhomogeneous Cosmological Models, Cambridge University Press, 1997.
- [7] T. Buchert, Gen. Rel. Gr. **40**, 467 (2008) gr-qc/0707.2153.
- [8] J. P. Boersma, Phys. Rev. **D57**, 798 (1998).